



Topics of limnological research in Mexico

Coordinator
Alfredo Pérez Morales

UNIVERSIDAD DE COLIMA

UNIVERSIDAD DE COLIMA

Dr. Christian Jorge Torres Ortiz Zermeño, Rector

Mtro. Joel Nino Jr., Secretario General

Mtro. Jorge Martínez Durán, Coordinador General de Comunicación Social

Mtro. Adolfo Álvarez González, Director General de Publicaciones

Mtra. Irma Leticia Bermúdez Aceves, Directora Editorial

Topics of limnological research in Mexico

Coordinator
Alfredo Pérez Morales



UNIVERSIDAD DE COLIMA

Topics of Limnological Research in Mexico

© UNIVERSIDAD DE COLIMA, 2025
Avenida Universidad 333
C.P 28040, Colima, Colima, México
Dirección General de Publicaciones
Telephone numbers: 312 316 1081 and 312 316 1000, extension: 35004
Email: publicaciones@ucol.mx
www.ucol.mx

ISBN electrónico: 978-968-9733-13-3
DOI: 10.53897/LI.2025.0032.UCOL
5E.1.1/317000/308/2024 Edición de publicación no periódica

All rights reserved according to law
Published in Mexico



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

You are free to: Share: copy and redistribute the material in any medium or format. Adapt: remix, transform, and build upon the material under the following terms: Attribution: You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use. NonCommercial: You may not use the material for commercial purposes. ShareAlike: If you remix, transform, or build upon the material, you must distribute your contributions under the same license as the original.

Cover and interior photographs: Juan Franco Rodríguez

Certified Publishing Process with ISO since 2005
Double blind ruling and editing are registered in the PRED Electronic Publishing System

Registration: LI-020-24
Received: September 2024
Dictamination: November 2024
Published: December 2025

*This book is dedicated to
Dr. Singaraju Sri Subrahmanyam Sarma,
in gratitude for all his teachings in the world of limnology.*



Index

Preface	10
Introduction	13
Analysis of the Ionic Quality of the Water in the North Aquifer and Cozumel Island, Quintana Roo, Mexico	16
<i>Gerardo Hernández-Flores, Martha Angélica Gutiérrez-Aguirre, Adrián Cervantes-Martínez.</i>	
Limnological Variations of a Tropical Semi-arid River Dam System, Central México	34
<i>Martín López-Hernández, Fernando González-Farías, María Guadalupe Ramos-Espinosa, Fernando Córdova-Tapia, Alejandro Gómez-Ponce.</i>	
Temporal Characterization of Water Quality of Rivers in Contrasting Zones of Two Watersheds in Veracruz, Mexico	58
<i>José Antolín Aké-Castillo, Miriam Guadalupe Ramos-Escobedo, Eduardo Aranda-Delgado.</i>	
Environmental Problems on Water Resources: A Review at the Basin Level with Emphasis on Tuxpan River in Veracruz, Mexico	77
<i>Blanca Esther Raya-Cruz, José Luis Alanís-Méndez, Carlos Francisco Rodríguez-Gómez, Karla Cirila Garcés-García.</i>	
Prospective Analysis of Major Phytoplankton Groups in Some Freshwater Bodies in Campeche, Southeastern Gulf of Mexico	94
<i>Juan Alfredo Gómez-Figueroa, Carlos Antonio Poot-Delgado, Jaime Rendón-von Osten, Yuri Okolodkov.</i>	

On the Relevance of Monitoring the Thermal Structure, Community Metabolism and Phytoplankton Ecology of Inland Waters of Mexico in the Context of Global Change	112
<i>Patricia Margarita Valdespino-Castillo, Jorge Alberto Ramírez-Zierold, Rocío Jetzabel Alcántara-Hernández, Mariel Barjau-Aguilar, Mario Alberto Neri-Guzmán, Paola Julieta Cortés Cruz, Oscar Alejandro Gerardo-Nieto, Martín Merino-Ibarra.</i>	
Middle-Term Hydrological and Microalgal Study in the Lower Basin of the Tuxpan River, Veracruz, Mexico	132
<i>Carlos Francisco Rodríguez-Gómez, Gabriela Vázquez, José Antolín Aké-Castillo, Angeles Rosseth Cruz-Ramírez.</i>	
Phytoplankton from two Dams in Central Mexico	153
<i>Gloria Garduño-Solórzano, José Manuel González-Fernández, Valeria Naomi Barranco-Vargas, Karla de la Luz-Vázquez, Cristian Alberto Espinosa-Rodríguez.</i>	
Towards Molecular, Genetic, and Optical Monitoring of Potentially Harmful Cyanobacteria Blooms in Mexican Freshwater Bodies	177
<i>Laura Valdés-Santiago, José Luis Castro-Guillén, Jorge Noé García-Chávez, Cynthia Paola Rangel-Chávez, Rosalba Alonso-Rodríguez, Alejandra Sarahí Ramírez-Segovia, Juan Gualberto Colli-Mull, Rafael Vargas-Bernal.</i>	
Free Living Continental Aquatic Ciliates (Alveolata: Ciliophora) from Mexico: An Overview of their Species Richness and Distribution	194
<i>Rosaura Mayén-Estrada, Carlos Alberto Durán-Ramírez, Fernando Olvera-Bautista, Víctor Manuel Romero-Niembro.</i>	
Potential Use of Rotifer and Cladoceran Diapausing Eggs as a Tool for Taxonomical, Ecological, and Evolutionary Studies	216
<i>Gerardo Guerrero-Jiménez, Elaine Aguilar-Nazare, Frida Sabine Álvarez-Solís, José Cristóbal Román-Reyes, Araceli Adabache-Ortiz, Marcelo Silva-Briano, Rocío Natalia Armas-Chávez.</i>	
Zooplankton Community and Trophic State in Lake Chapala	234
<i>Cristian Alberto Espinosa-Rodríguez, Lizbeth Cano-Parra, Omar Alfredo Barrera-Moreno.</i>	

Seasonal and Diel Influence of Environmental Factors on the Parameters of a Zooplankton Community in a Tropical Coastal Lagoon	255
<i>Manuel Castillo-Rivera.</i>	
Utilization of Zooplankton in Environmental Risk Assessment in Mexico	275
<i>Cesar Alejandro Zamora-Barrios, Rosa Martha Moreno-Gutiérrez,</i>	
<i>Uriel Arreguin-Rebolledo, Mario Joshue Espinosa-Hernández,</i>	
<i>Francisco José Torner-Morales.</i>	
Exploring Zooplankton-Macrophytes Interaction Research in Mexico: Bibliometric Analysis	296
<i>Marco Antonio Jiménez-Santos, Michael Anai Figueroa-Sánchez.</i>	
The Freshwater and Brackish Hydrozoans of Mexico: An Overview of their Diversity	315
<i>José María Ahuatzin-Hernández, Lorena Violeta León-Deniz.</i>	
Aquatic Macroinvertebrates Diversity in the Grijalva and Usumacinta Rivers, Mexico	332
<i>Everardo Barba-Macías, Juan Juárez-Flores, Cinthia Trinidad-Ocaña,</i>	
<i>José Francisco Miranda-Vidal.</i>	
Fishing Among Socioecological Challenges: The Case of the Zimapán Dam	361
<i>Brenda Rodríguez-Cortés, Karina E. Ruiz-Venegas, Martín López-Hernández,</i>	
<i>Alejandro Gómez-Ponce, Fernando Córdova-Tapia.</i>	
Conclusions	379
About the authors	381
Acknowledgements	395



Seasonal and Diel Influence of Environmental Factors on the Parameters of a Zooplankton Community in a Tropical Coastal Lagoon

Manuel Castillo-Rivera

Laboratorio de Peces, Departamento de Biología, Universidad Autónoma Metropolitana, Iztapalapa. Av. Ferrocarril San Rafael Atlixco, Núm. 186, Col. Leyes de Reforma 1a Sección, Alcaldía Iztapalapa, Mexico City 09310, Mexico. E-mail: crma@xanum.uam.mx

Abstract

Zooplankton communities show seasonal and diel changes mainly associated with the variability of abiotic and biotic factors. Community parameters are good descriptors for evaluating these changes. The study's objective was to analyze the seasonal and diel variation of the parameters of a zooplanktonic community in the La Mancha Lagoon (Veracruz state, Mexico), and the influence exerted by environmental factors. For a year, a 24-h cycle was completed each month, sampling zooplankton every 4 h during a new moon. Abiotic and biotic variables were recorded simultaneously. Univariate permutational analysis of variance, canonical correspondence analysis, and simple correlation analysis were used to analyze the data. Species richness, diversity, and dominance showed significant differences at the seasonal and diel levels. Seasonally, diversity and species richness showed their highest values in November-December and April, and low values in May, while dominance presented a completely inverse pattern. At the diel level, species richness and diversity showed high values at night, while dominance at dawn. Diversity was strongly associated with dominance inversely and directly correlated with species richness. Canonical correspondence ordination explained a high percentage of the constrained variance of the parameters-environment relationship (99.2 %), revealing salinity and diel effect as the most critical variables driving community structure. The influence of salinity was observed at a monthly and daily level. A negligible effect of biotic variables (chlorophyll a and densities of diatoms, dinoflagellates, and cyanobacteria) was also observed. Due to the high environmental variability in estuaries, abiotic factors may play a more important role in community structuring.

Keywords

Abiotic factors, environmental variability, dominant species, predator-evasion, constrained ordination.

Introduction

Zooplankton communities are of great importance in freshwater, brackish, and marine systems, playing an essential role in energy transfer in the aquatic food web between primary producers and higher consumers and contributing to nutrient recycling (Dvoretsky & Dvoretsky, 2021; Muñoz-Colmenares et al., 2021; Rosa et al., 2021). Furthermore, zooplanktonic organisms are susceptible to environmental changes. They are therefore considered good indicators of ecosystems, because a change in abiotic and biotic factors in aquatic systems results in a change in the relative composition and abundance of these organisms (Azevêdo et al., 2015; Arias et al., 2022; Guermazi et al., 2023).

An objective of aquatic ecological research is to know the response of planktonic communities to environmental conditions (Marques et al., 2009). Thus, many studies have demonstrated seasonal and diel changes in the species composition of zooplanktonic communities. Seasonally, these changes have been explained in terms of the influence of abiotic factors, such as salinity, temperature, dissolved oxygen, and light (Shi et al., 2015; Ge et al., 2021; Ursella et al., 2021; Arias et al., 2022), and/or by biotic factors, such as phytoplankton abundance, chlorophyll a, predation, and competition (Węgleńska et al., 1997; García-Herrera et al., 2022), which can act separately or in synergy (Liang et al., 2020; Hobbs et al., 2021; Muñoz-Colmenares et al., 2021; Guermazi et al., 2023).

Diel changes are associated with vertical migration, which is a ubiquitous characteristic in marine and freshwater planktonic communities (van Haren & Compton, 2013; Ursella et al., 2021), and these have been mainly related to food availability, predation avoidance and competition, and are generally modulated by diel and tidal cycles (Hobbs et al., 2021; García-Herrera et al., 2022). Indeed, several studies have observed a considerable increase in the number of species during the night (Marques et al., 2009; Primo et al., 2012) and for this reason, samplings throughout 24 h cycles can provide a better understanding of zooplankton richness.

Although the study of zooplankton communities is a multivariate process, because many species are captured in several samples, there are good univariate descriptors that allow condensing and summarizing information about the number, identity, and relative abundance of the species and, consequently, analyze changes in the community structure. These descriptors are community properties or community parameters, which are essentially species richness, diversity, evenness, and dominance (Morin, 2011), which ha-

ve been used in ecological studies of zooplankton communities (Liang et al., 2020; Arias et al., 2022; Romero et al., 2022; Taniguchi et al., 2023).

There are many studies on the structure of zooplankton communities in freshwater, estuarine, and marine systems. Still, in the particular case of Mexico, there are few studies that simultaneously address seasonal and diel changes in the structure of these communities. In this sense, the main objective of this study was to analyze the seasonal and diel variation of richness, diversity, evenness, and dominance of a zooplanktonic community in La Mancha Lagoon, and the influence exerted by different environmental factors, considering the hypothesis that community structure will be affected mainly by seasonal variability of abiotic factors and diel variation mainly by biotic factors.

Materials and Methods

Study Area

La Mancha Lagoon is a Ramsar site, a category defined as a site that provides for national action and international cooperation in the conservation of wetlands and the rational and sustainable use of their resources. This system is located in the state of Veracruz, Mexico ($19^{\circ}33'55''$ – $19^{\circ}35'44''$ N and $96^{\circ}22'41''$ – $96^{\circ}23'39''$ W). It is a small intermittent microtidal system with an approximate extension of $1,742 \text{ km}^2$ (Morgado-Dueñas & Castillo-Rivera, 2022). The region has a warm, sub-humid climate (Köppen climate classification: Aw2), with two climatic seasons: a dry season (mean monthly rainfall $<60 \text{ mm}$) from November to May and a rainy season (mean monthly rainfall $>100 \text{ mm}$) from June to October.




Figure 1. Geographical Location of the the La Mancha Lagoon.

Sampling Collections

For a year, the zooplanktonic community at the mouth of La Mancha Lagoon was sampled through monthly diel cycles. Thus, 24-h cycle was completed each month, taking samples every four hours (6 samples per cycle) during the new moon period to maximize the light/dark effect (Castillo-Rivera et al., 2010). Zooplankton samples were obtained by circular surface hauls using a standard net (100 cm length, 30 cm diameter, and 150 µm mesh size). Organisms were immediately preserved in 4 % borax-buffered formalin. Simultaneously, the biotic factors chlorophyll a (spectrophotometric method, SCOR, 1966), and abundance of diatoms, dinoflagellates, and cyanobacteria (Utermöhl method, using an inverted microscope Motic AE31), and the abiotic variables salinity (ATAGO S-10E Refractometer), temperature, dissolved oxygen (YSI 550A DO Instrument), water level (tidal phase), and time of day were recorded, according to Mecalco-Hernández et al. (2018). For quantitative analysis, the time of day was classified into diel periods, considering a light-dark gradient on an ordinal scale with values of 2 (day=10:00 and 14:00 h), 1 (twilight=06:00 and 18:00 h) and 0 (night=22:00 and 2:00 h).

Data Analysis

Fifty-seven zooplankton components were recorded, corresponding to 55 species, in addition to two undetermined stages (zoea larvae of brachyura and fish eggs). Their abundances were expressed as individual numbers per cubic meter (ind/m³) (Mecalco-Hernández et al., 2018; Mecalco-Hernández & Castillo-Rivera, 2020). From these data, community parameters were estimated for each sample: species richness (number of species), diversity (Shannon-Winer index), evenness (Pielou index), and dominance (Simpson index).

Two-way univariate permutational analysis of variance (PERMANOVA) was applied to evaluate significant differences in the analyzed parameters among months and diel periods, as well as their interactions. PERMANOVA is a routine for testing the response of one or more variables to one or more factors based on a resemblance measure. This method is also appropriate because it uses multiple random permutations to obtain *p*-values; thus, the permutation procedure directly implies normality and homogeneity of variances (Anderson, 2001; Anderson et al., 2008). PERMANOVA was performed using Euclidean distance and permuted residuals under a reduced model, Type III (maximum permutations=999), according to the routine for univariate analysis (Anderson et al., 2008). This analysis was performed using the software PRIMER v7.

To evaluate the influence of environmental factors on community structure, canonical correspondence analysis (CCA) was applied to the parameter data matrix (dependent set: community parameters) and the factor data matrix (explanatory set: abiotic and biotic variables). Inter-set correlations from this analysis were used to determine the environmental

variables that were most important in determining the variability of community parameters (McGarigal et al., 2000). The significance of each of these factors was determined using 499 unrestricted Monte Carlo permutations. A biplot of parameters and explanatory factors was constructed to observe any pattern associated with these factors. All of these analyses were performed using the package CANOCO ver. 4.5 (ter Braak & Šmilauer, 2002). In addition, the degree of simple association between two variables was evaluated using non-parametric Spearman correlation analysis (r_s).

Results

The seasonal pattern of zooplankton diversity and species richness showed high values in November-December and April, and low values in May, July, and January (Fig. 2A). In contrast, dominance was high in May (defined by the copepods *Tortanus (Acutanus) setacaudatus* and *Pseudosiacptomus pelagicus*), October (by zoea larvae of brachyura), and March (by the copepod *Acartia (Acanthacartia) tonsa*), with low values in December and April, while evenness shows a completely inverse pattern to dominance (Fig. 2B).




Figure 2. Monthly Variation of: A) Mean and Standard Error Values of diversity and species richness, and B) Mean and Standard Error Values of Dominance and Evenness (for each Monthly Value n=6).

The diel variability of these parameters showed that diversity and richness presented high values at night (22:00 to 02:00 h) and low values at dawn (06:00 h) or dusk (18:00 h) and during the day (10:00 to 14:00 h) (Fig. 3A). On the contrary, the high value of dominance occurred at dawn (06:00 h) dominated by *A. tonsa* and zoea larvae, and low values during the night (22:00 to 02:00 h) while the evenness shows a reverse pattern (Fig. 3B). The community was dominated solely by the abundance of four zooplanktonic compo-

nents, *A. tonsa*, *P. pelagicus*, *T. setacuadatus*, and zoea larvae, which represented 89 % of the total abundance in number of the entire community.




Figure 3. Mean and Standard Error Values by Hour: A) Diversity and Species Richness, and B) Dominance and Evenness (for each Hourly Value n=12).

According to the two-way univariate results (PERMANOVA), all parameters show significant differences among months. Similarly, at the diel level, there were significant differences except for evenness. All interactions between monthly and diel factors were not significant (P's >0.3, Table 1).

Table 1. Results of Two-way Univariate Analysis of Variance (PERMANOVA) to Test the Effects of the Month and Diel Periods (Day, Twilight, and Night) on the Community Parameters.

Source of Variance	df	MS	Pseudo-F	P (Permuted)
Diversity				
Months	11	2.589	6.053	0.001
Diel	2	1.563	3.655	0.039
Months x Diel	22	0.371	0.866	0.638
Error	32	0.428		
Species Richness				
Months	11	146.19	5.822	0.001
Diel	2	232.96	9.278	0.001
Months x Diel	22	16.320	0.650	0.835
Error	32	25.109		
Dominance				
Months	11	0.178	4.455	0.001
Diel	2	0.179	4.479	0.020
Months x Diel	22	0.044	1.096	0.396
Error	32	0.040		
Evenness				
Months	11	0.127	5.019	0.001
Diel	2	0.041	1.627	0.246
Months x Diel	22	0.030	1.178	0.328
Error	32	0.025		

In this sense, diversity was mainly inversely correlated with dominance ($r_s=-0.981$, $P<0.0001$) and directly correlated with species richness ($r_s 0.747$, $P<0.0001$), while species richness and dominance were also negatively correlated ($r_s=-0.693$, $P<0.0001$).

Concerning environmental factors' influence on community structure, the simultaneous ordination of parameters and environmental matrices in CCA showed that all canonical axes were significant ($P=0.010$). The first two axes explained 99.2 % (76.9 and 22.3 %, respectively) of the total constrained variance. Intra-set correlations indicated that salinity, the diel effect, and temperature were the most critical variables in structuring the zooplankton community. Likewise, according to Monte Carlo significance tests, only these factors significantly affected the community parameters. Both inter-set correlations and significance tests indica-

ted that biotic factors such as chlorophyll a and the abundance of diatoms, dinoflagellates, and cyanobacteria were unimportant in community structuring (Table 2).

Table 2. Inter-set Correlations Between Environmental Factors and Species Scores from the Canonical Correspondence Analysis. The Significance of these Factors (Unrestricted Monte Carlo Permutations) Is also Shown.

Environmental Factors	Inter-set Correlations		
	Axis 1	Axis 2	P(Permuted)
Salinity	-0.510	0.104	0.002
Diel Effect	0.363	0.052	0.020
Temperature	-0.072	-0.284	0.038
Tidal Stage	-0.149	-0.156	0.192
Dissolved Oxygen	0.133	-0.145	0.272
Diatoms	-0.129	-0.066	0.506
Chlorophyll a	0.199	0.068	0.608
Cyanobacteria	0.181	-0.105	0.748
Dinoflagellates	0.059	-0.023	0.934

In the CCA analysis biplot (Fig. 4), the length and direction of the arrows indicate the relative importance and direction of change that each factor has in the ordination. This diagram reveals a main ordination gradient related to salinity that is directly associated with species richness and inversely with dominance. Another major trend, perpendicular to the first, is related to temperature and is in the opposite direction of the diel effect, which shows that diversity is associated with low temperatures and evenness in the daytime.

Thus, salinity was the environmental variable that was most correlated individually with the community parameters, directly with diversity ($rs=0.575$, $P<0.0001$), species richness ($rs=0.491$, $P<0.0001$), and evenness ($rs=0.486$, $P=0.0001$), and inversely with dominance ($rs=-0.536$, $P<0.0001$). The direct relationship between salinity, diversity, and species richness can be observed at the seasonal and diel levels (Fig. 5A-B and 5C-D, respectively). Dominance has an inverse clear relationship with salinity only seasonally. Other abiotic and biotic variables did not show significant simple correlations with the parameters.





Figure 4. Bi-plot of Canonical Correspondence Analysis of the Overall Community Parameters and Environmental Variable Data Matrices. Data Points Represent Parameters (\blacktriangle), and Arrows Represent the Direction and Relative Importance of Explanatory Variables.



Continued from previous page.




Figure 5. Monthly Mean Values of Salinity and Community Parameters: A) Direct Relationship between Salinity (●), Species Richness (■), and Diversity (▲), and B) Inverse Relationship between Salinity (●) and Dominance (▲) (for Mean Monthly Values $n=6$). Mean Values by Hour of Salinity and Community Parameters: C) Direct Relationship between Salinity (●), Species Richness (■), and Diversity (▲), and D) Relationship between Salinity (●) and Dominance (▲) (for Mean Hourly Values $n=12$).

Discussion

Although the formal study of zooplankton in Mexican waters began more than a century ago, most of the studies carried out correspond mainly to research in freshwater and marine systems with central reference only to some groups of zooplankton (Mecalco-Her-

nández & Castillo-Rivera, 2020; Alcocer et al., 2022; Cervantes-Martínez et al., 2023), there are few studies in estuarine systems, especially in the state of Veracruz. Likewise, few studies have inferred diel changes in zooplankton communities from low-frequency temporal sampling which allows a more precise assessment of the timing of diel changes throughout the 24 h cycle and the influence of environmental factors on these changes over short periods.

In the research of zooplankton communities, many studies have used community parameters to describe changes in community structure adequately (Liang et al., 2020; Arias et al., 2022; Romero et al., 2022; Taniguchi et al., 2023). In the present study, all community parameters showed a significant monthly variation, which is related to the strong environmental variability that characterizes estuarine ecosystems and to the fact that zooplankton usually presents a relatively rapid response to environmental changes (Benfield, 2013; Guermazi et al., 2023).

In this way, diversity and species richness showed pulses in December and April and low values in May, while dominance showed a completely inverse pattern. Indeed, simple correlations showed that diversity and species richness had a strong inverse correlation with dominance. Concerning environmental stability, diversity can be determined primarily by evenness in habitats with low variability. However, in environments with strong variability, such as estuaries, few eurytopic species (tolerant to a wide range of ecological conditions) would be expected to be dominant, so diversity would mainly be inversely affected by dominance. Indeed, in La Mancha Lagoon, almost 90 % of the numerical abundance of the entire community was dominated by only four zooplankton components (*A. tonsa*, *P. pelagicus*, *T. setacudatus* and zoea larvae).

At the diel level, all parameters showed significant differences among hours, except evenness. Species richness and diversity showed high values during the night and low values during the day. According to the PERMANOVA results, there was no significant interaction between the month and diel period factors, for all four parameters, indicating that the diel pattern was consistent across all sampled months. In this way, many studies have indicated that during the night there is a significant increase in the number of species (Marques et al., 2009; Primo et al., 2012). This behavioral pattern has been considered the most common characteristic in marine and freshwater planktonic communities, which is characterized by an evening ascent and a morning descent (van Haren & Compton, 2013; Ursella et al., 2021).

Considering the predator avoidance hypothesis, the vertical migration of zooplankton to deeper waters during daylight hours, where the probability of being detected by visually searching predators is lower than if they remained in shallow waters with more illuminated conditions, and during the night, under dark conditions, zooplankton migrate

upward to feed and shelter (Węgleńska et al., 1997; Ursella et al., 2021; Garcia-Herrera et al., 2022). These vertical patterns may also appear to be influenced by abiotic factors such as hydrological regime (inputs of freshwater flow), water currents, light variation, tidal phase, and salinity (Primo et al., 2012; Benfield, 2013; Hobbs et al., 2021; Ge et al., 2021).

The dominant zooplanktonic components, *A. tonsa* and zoea larvae, showed a typical crepuscular occurrence with pulses at dawn. In this sense, the increase in activity during twilight hours may be related to compensation strategies that attempt to resolve the trade-off between feeding and avoiding predators. Twilight conditions may mitigate visual detection by visual predators, but provide sufficient light to detect prey. Indeed, considering a predator-prey encounter model (Giske et al., 1994), for smaller zooplankters the risk of mortality will generally be reduced in accordance with a general decrease in ambient light.

Regarding the knowledge of biodiversity in zooplankton communities, studies over short periods throughout sampling in 24 h cycles can provide a better understanding of zooplankton richness. Indeed, in the present study, of the total 57 zooplanktonic components recorded, 13 (~23 %) were collected only at twilight and during the night.

About the multivariate influence of environmental factors on community parameters, the CCA ordination explained a high percentage of the constrained variance of the parameters-environment relationship (99.2 %), which implies that the environmental variables considered give an adequate description of the community structure. According to inter-set correlations and Monte Carlo significance tests, abiotic factors salinity and diel effect (related to light gradient) were the most critical variables driving community structure.

Indeed, salinity is one of the main factors driving the structuring of zooplankton communities in estuarine and marine systems (Benfield, 2013; Shi et al., 2015; Dvoretsky & Dvoretsky, 2017; Chen et al., 2023), affecting the richness and diversity species inversely (negative correlations) or directly (positive correlations), depending on whether the communities are dominated by freshwater or marine species (Paturej & Gutkowska, 2015; Helenius et al., 2017; Yuan et al., 2020).

In the present study, according to CCA results salinity has a positive effect on species richness, while it has a negative effect on dominance. These results were also confirmed by simple correlations, which indicated significant direct correlations of salinity with species richness and diversity and a significant inverse correlation between salinity and dominance. Higher salinities allowed the entry of a greater number of neritic species to the lagoon, thus increasing richness and diversity. In contrast, in low-salinity conditions only tolerant species remain dominant in the community. Thus, the effect of salinity was observed at monthly and diel levels. The light gradient showed that species richness was more significant during the night, while other abiotic factors such as dissolved oxygen and tidal stage showed little incidence on the community structure (low inter-set correlations and no

significant effect). This is because the dissolved oxygen values in the system are probably greater than the limits of lethal and sublethal hypoxic conditions. Likewise, although some studies have observed the importance of the tide in the structuring of the community (Marques et al., 2009; Primo et al., 2012), for the zooplanktonic community of La Macha, the tide stage had little influence mainly because the southwest coast of the Gulf of Mexico is predominantly microtidal (Ellis & Dean, 2012), with <20 cm within the lagoon.

CCA results also showed the negligible effect of biotic factors on community structuring. Many studies have observed that zooplankton communities can be affected by biotic factors such as phytoplankton density (García-Herrera et al., 2022; Guermazi et al., 2023) and chlorophyll a (Liang et al., 2020; Muñoz-Colmenares et al., 2021; Chen et al., 2023), but in the present study, chlorophyll a and the densities of diatoms, dinoflagellates, and cyanobacteria were not significant. Thus, both abiotic and biotic conditions can have an important incidence in a structuring community, but their relative importance may vary among different environmental conditions and across temporal and spatial scales. While abiotic factors can affect community structure by setting the limits of abiotic tolerance of species, biotic factors may also affect diversity patterns by influencing species abundance and composition (Måsviken et al., 2023).

In this way, many studies have compared the relative effect of these factors on zooplankton communities, observing that both can act synergistically (Gray et al., 2012; Mecalco-Hernández et al., 2018; Yang et al., 2019; Chará-Serna and Casper, 2021) or at a different temporal scale (Gabaldón et al., 2019; Rollwagen-Bollens et al., 2020). In this way, the variability of abiotic factors can also modify the biological response of species (Begon & Townsend, 2021), as observed in La Mancha Lagoon, where the light/dark cycle affects predator-evasion patterns.

Notwithstanding the above, it would be expected that in environments with low environmental variability (stable), biotic factors such as food availability, predation, and competition would be the main regulators of the community. In contrast, in environments with high environmental variability, abiotic factors would play a more important role, as is also observed in the zooplanktonic community of La Mancha Lagoon and other similar systems (Sahuquillo & Miracle, 2019; Sgarzi et al., 2019).

Conclusion

A better understanding of the zooplanktonic community dynamics of aquatic ecosystems can be achieved by sampling over 24 h. Thus, species richness was consistently greater during the night, which was related to predator-evasion strategies modulated by the light/dark cycle. The environmental variables considered in the present study explained a very high percentage of the community structure's temporal variability, allowing a more reliable interpretation of the results. The high environmental variability characteristic of estuaries caused the community structure to vary significantly among months and hours. As a consequence of this variability, diversity was mainly associated with dominance, and the abiotic factors salinity and diel effect played a more important role in determining the community structure, while the biotic variables chlorophyll a, and densities of diatoms, dinoflagellates, and cyanobacteria had negligible importance.

Acknowledgements

This research was funded by Universidad Autónoma Metropolitana, Unidad Iztapalapa (Project: 14302047). I thank Ángeles Mecalco-Hernández for her valuable help in data processing. Thanks also to the editor and the constructive comments on the manuscript from two anonymous reviewers, which were very valuable.

References

Alcocer, J., Espinosa-Rodríguez, C. A., Fernández, R., Lugo-Vázquez, A., Macek, M., Maeda-Martínez, A. M., Martínez-Jerónimo, F., Ortega-Mayagoitia, E., & Oseguera, L. A. (2022). The ecology of the zooplankton in Mexican inland waters: What we know so far. *Limnologica*, 9, 126037. <https://doi.org/10.1016/j.limno.2022.126037>.

Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance. *Austral Ecology*, 26, 32-46. <https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x>

Anderson, M. J., & Gorley, R. N., [https://doi.org/Clarke, K. R. \(2008\). PERMANOVA + for PRIMER: Guide to Software and Statistical Methods. Massey University, Palmerston North.](https://doi.org/Clarke, K. R. (2008). PERMANOVA + for PRIMER: Guide to Software and Statistical Methods. Massey University, Palmerston North.)

Arias, M. J., Vaschetto, P. A., Marchese, M., Regaldo, L., & Gagneten, A. M. (2022). Benthic macroinvertebrates and zooplankton communities as ecological indicators in urban wetlands of Argentina. *Sustainability*, 14, 4045. <https://doi.org/10.3390/su14074045>

Azevêdo, D. J. S., Barbosa, J. E. L., Gomes, W. I. A., Porto, D. E., Marques, J. C., & Molozzi, J. (2015). Diversity measures in macroinvertebrate and zooplankton communities related to the trophic status of subtropical reservoirs: Contradictory or complementary responses? *Ecological Indicators*, 50, 135-149. <https://doi.org/10.1016/j.ecolind.2014.10.010>

Begon, M., & Townsend, C. R. (2021). *Ecology: From Individuals to Ecosystems* (5th ed.). John Wiley & Sons.

Benfield, M. C. (2013). Estuarine Zooplankton. In Day, J. W., Crump, B. C., Kemp, W. M., Yáñez-Arancibia, A. (Eds.) *Estuarine ecology*. Wiley-Blackwell. <https://doi.org/10.1002/9781118412787.ch11>

Castillo-Rivera, M., Zárate-Hernández, R., Ortiz-Burgos, S., & Zavala-Hurtado, A. (2010). Diel and seasonal variability in the fish community structure of a mud-bottom estuarine habitat in the Gulf of Mexico. *Marine Ecology*, 31(4), 633-642. <https://doi.org/10.1111/j.1439-0485.2010.00394.x>

Chará-Serna, A. M., & Casper, A. (2021) How do large river zooplankton communities respond to abiotic and biotic drivers over time? A complex and spatially dependent example. *Freshwater Biology*, 66(2), 391-405. <https://doi.org/10.1111/fwb.13646>

Cervantes-Martínez, A., Durán Ramírez, C. A., Elías-Gutiérrez, M., García-Morales, A. E., Gutiérrez-Aguirre, M., Jaime, S., Macek, M., Maeda-Martínez, A. M., Martínez-Jerónimo, F., Mayén-Estrada, R., Medina-Durán, J. H., Montes-Ortiz, L., Olvera-Bautista, J. F. Y., Romero-Niembro, V. M., & Suárez-Morales, E. (2023). Freshwater diversity of zooplankton from Mexico: Historical review of some of the main groups. *Water*, 15, 858. <https://doi.org/10.3390/w15050858>

Chen, X., Cui, Z., Zhang, Y., Zhang, X., Chen, J., & Wei, Y. (2023). Contrasting effects of river inflow and seawater intrusion on zooplankton community structure in Jiaozhou bay, the Yellow Sea. *Marine Environmental Research*, 192, 106194. <https://doi.org/10.1016/j.marenvres.2023.106194>.

Dvoretsky, V. G., & Dvoretsky, A. G. (2017). Macrozooplankton of the Arctic – The Kara Sea in relation to environmental conditions. *Estuarine, Coastal and Shelf Science*, 188, 38-55. <https://doi.org/10.1016/j.ecss.2017.02.008>.

Dvoretsky, V. G., & Dvoretsky, A. G. (2021). Winter zooplankton in a small Arctic lake: abundance and vertical distribution. *Water*, 13, 912. <https://doi.org/10.3390/w13070912>

Ellis, J. T., & Dean, B. J. (2012). Gulf of Mexico Processes. *Journal of Coastal Research*, (Special Issue 60), 6-13. https://doi.org/10.2112/SI_60_2

Gabaldón, C., Devetter, M., Hejzlar, J., Simek, K., Znachor, P., Nedoma, J., & Sed'a, J. (2019). Seasonal strengths of the abiotic and biotic drivers of a zooplankton community. *Freshwater Biology*, 64(7), 1326–1341. <https://doi.org/10.1111/fwb.13308>

Garcia-Herrera, N., Cornils, A., Laudien, J., Niehoff, B., Höfer, J., Försterra, G., González, H. E., & Richter, C. (2022). Seasonal and diel variations in the vertical distribution, composition, abundance and biomass of zooplankton in a deep Chilean Patagonian Fjord. *PeerJ*, 10, e12823. <https://doi.org/10.7717/peerj.12823>

Ge, R., Chen, H., Liu, G., Zhu, Y., & Jiang, Q. (2021). Diel vertical migration of mesozooplankton in the northern Yellow Sea. *Journal of Oceanology and Limnology*, 39(4), 1373-1386. <https://doi.org/10.1007/s00343-020-0170-y>

Giske, J., Aksnes, D. L., & Fiksen, Ø. (1994). Visual predators, environmental variables and zooplankton mortality risk. *Vie et Milieu*, 44(1), 1-9. <https://hal.sorbonne-universite.fr/hal-03047873>

Gray, D. K., Arnott, S. E., Shead, J. A., & Derry, A. M. (2012). The recovery of acid-damaged zooplankton communities in Canadian Lakes: the relative importance of abiotic, biotic and spatial variables. *Freshwater Biology*, 57(4), 741-758. <https://doi.org/10.1111/j.1365-2427.2012.02739.x>

Guermazi, W., El-khateeb, M., Abu-Dalo, M., Sallemi, I., Al-Rahahleh, B., Rekik, A., Belmonte, G., Ayadi, H., & Annabi-Trabelsi, N. (2023). Assessment of the zooplankton community and

water quality in an artificial Freshwater lake from a semi-arid area (Irbid, Jordan). *Water*, 15, 2796. <https://doi.org/10.3390/w15152796>

Helenius, L. K., Leskinen, E., Lehtonen, H., & Nurminen, L. (2017). Spatial patterns of littoral zooplankton assemblages along a salinity gradient in a brackish sea: A functional diversity perspective. *Estuarine, Coastal and Shelf Science*, 198(Part B), 400-412. <https://doi.org/10.1016/j.ecss.2016.08.031>.

Hobbs, L., Banas, N. S., Cohen, J. H., Cottier, F. R., Berge, J., & Varpe, Ø. (2021). A marine zooplankton community vertically structured by light across diel to interannual timescales. *Biology Letters*, 17, 20200810. <https://doi.org/10.1098/rsbl.2020.0810>

Liang, D., Wang, Q., Wei, N., Tang, C., Sun, X., & Yang, Y. (2020). Biological indicators of ecological quality in typical urban river-lake ecosystems: The planktonic rotifer community and its response to environmental factors. *Ecological Indicators*, 112, 106127. <https://doi.org/10.1016/j.ecolind.2020.106127>

Marques, S. A. Azeiteiro, U. M., Martinho, F., Viegas, I., & Pardal, M. A. (2009). Evaluation of estuarine mesozooplankton dynamics at a fine temporal scale: the role of seasonal, lunar and diel cycles. *Journal of Plankton Research*, 31(10), 1249-1263. <https://doi.org/10.1093/plankt/fbp068>

Måsviken, J., Dalén, L., Norén, K., & Dalerum, F. (2023). The relative importance of abiotic and biotic environmental conditions for taxonomic, phylogenetic, and functional diversity of spiders across spatial scales. *Oecologia*, 202(2), 261-273. <https://doi.org/10.1007/s00442-023-05383-0>

McGarigal, K., Cushman, S., & Stafford, S. (2000). *Multivariate Statistics for Wildlife and Ecology Research*. Springer-Verlag. <https://doi.org/10.1007/978-1-4612-1288-1>

Mecalco-Hernández, Á., & Castillo-Rivera, M. (2020). Riqueza zooplanctónica en la boca de la Laguna La Mancha, durante muestreos mensuales y nictímeros. *Hidrobiológica*, 30(2), 143-153. <https://doi.org/10.24275/uam/izt/dcbs/hidro/2020v30n2/Castillo>

Mecalco-Hernández, A., Castillo-Rivera, M., Sanvicente-Añorve, L., Flores-Coto, C., & Álvarez-Silva, C. (2018). Variación estacional y nictímera en la distribución del zooplancton dominante en una laguna costera tropical. *Revista de Biología Marina y Oceanografía*, 53, 39-49. <https://doi.org/10.4067/S0718-19572018000100039>

Morgado-Dueñas, G., & Castillo-Rivera, M. (2022). Diversity, seasonal and diel distribution patterns of Anchovies (Osteichthyes) in a protected tropical lagoon in the Southwestern Gulf of Mexico. *Diversity*, 14, 852. <https://doi.org/10.3390/d14100852>

Morin, J. P. (2011). *Community Ecology*. (2nd ed.). Wiley-Blackwell. <https://doi.org/10.1002/9781444341966>

Muñoz-Colmenares, M. E., Vicente, E., Soria J. M., & Miracle, M. R. (2021). Zooplankton changes at six reservoirs in the Ebro watershed, Spain. *Limnetica*, 40(2), 279-294. <https://doi.org/10.23818/limn.40.19>

Paturej, E., & Gutkowska, A. (2015). The effect of salinity levels on the structure of zooplankton communities. *Archives of Biological Sciences, Belgrade*, 67(2), 483-492. <https://doi.org/10.2298/ABS140910012P>

Primo, A. L., Azeiteiro, U. M., Marques, S. C., Ré, P., & Pardal, M. A. (2012). Vertical patterns of ichthyoplankton at the interface between a temperate estuary and adjacent coastal waters: Seasonal relation to diel and tidal cycles. *Journal of Marine Systems*, 95, 16-23. <https://doi.org/10.1016/j.jmarsys.2011.12.008>

Romero, N., Attademo, A. M., Reno, R., Regaldo, L., Repetti, R. M., & Gagneten, A. M. (2022). Analysis of the zooplanktonic community in rice fields during a crop cycle in agroecological versus conventional management. *Limnetica*, 41(1), 107-120. <https://doi.org/10.23818/limn.41.09>

Rosa, L. M., Cardoso, L. S., Rodrígues, L. R., & Motta-Marques, D. (2021). Density versus biomass responses of zooplankton to environmental variability in a subtropical shallow lake. *Inland Waters*, 11(1), 44-56. <https://doi.org/10.1080/20442041.2020.1714383>

Rollwagen-Bollens, G., Bollens, S., Dexter, E., & Cordell, J. (2020). Biotic vs. abiotic forcing on plankton assemblages varies with season and size class in a large temperate estuary. *Journal of Plankton Research*, 42(2), 221-237. <https://doi.org/10.1093/plankt/fbaa010>

Sahuquillo, M., & Miracle, R. M. (2019). Rotifer communities in Mediterranean ponds in eastern Iberian Peninsula: abiotic and biotic factors defining pond types. *Limnetica*, 38(1), 103-117. <https://doi.org/10.23818/limn.38.14>

Sgarzi, S., Badosa, A., Leiva-Presa, À., Benejam, L., López-Flores, R., & Brucet, S. (2019). Plankton taxonomic and size diversity of Mediterranean brackish ponds in spring: influence of abiotic and biotic factors. *Water*, 11(1), 106. <https://doi.org/10.3390/w11010106>

Scientific Committee on Oceanic Research (SCOR). (1966). *Determination of photosynthetic pigments in seawater. Monographs on Oceanographic Methodology*. UNESCO.

Shi, Y.-Q., Song, S., Zhang, G.-T., Wang, S.-W., & Li, C.-L. (2015). Distribution pattern of zooplankton functional groups in the Yellow Sea in June: a possible cause for geographical separation of giant jellyfish species. *Hydrobiologia*, 754, 43-58. <https://doi.org/10.1007/s10750-014-2070-7>

Taniguchi, R., Amei, K., Tokuhiro, K., Yamada, Y., Kitamura, M., & Yamaguchi, A. (2023). Diel, seasonal and vertical changes in the pelagic amphipod communities in the subarctic Pacific: insights from imaging analysis. *Journal of Plankton Research*, 45(3), 554-570. <https://doi.org/10.1093/plankt/fbad017>

ter Braak, C. J. F., & Šmilauer, P. (2002). *CANOCO reference manual and CanoDraw for Windows. User's guide: software for canonical community ordination* (version 4.5). www.canoco.com.

Ursella, L., Pensieri, S., Pallàs-Sanz, E., Herzka, S. Z., Bozzano, R., Tenreiro, M., Cardin, V., Candela, J., & Sheinbaum, J. (2021). Diel, lunar and seasonal vertical migration in the deep western Gulf of Mexico evidenced from a long-term data series of acoustic backscatter. *Progress in Oceanography*, 195, 102562. <https://doi.org/10.1016/j.pocean.2021.102562>.


Van Haren, H., & Compton, T. J. (2013). Diel vertical migration in deep sea plankton is finely tuned to latitudinal and seasonal day length. *PLoS ONE*, 8(5), e64435. <https://doi.org/10.1371/journal.pone.0064435>

Węgleńska, T., Ejsmont-Karabin, J., & Rybak, J. I. (1997). Biotic interactions of the zooplankton community of a shallow, humic lake. *Hydrobiologia*, 342, 185-195. <https://doi.org/10.1023/A:1017062602104>

Yang, Y., Gao, Y., Chen, Y., Li, S., & Zhan, A. (2019). Interactome-based abiotic and biotic impacts on biodiversity of plankton communities in disturbed wetlands. *Diversity and Distributions*, 25(9), 1416-1428. <https://doi.org/10.1111/ddi.12949>

Yuan, D., Chen, L., Luan, L., Wang, Q., & Yang, Y. (2020). Effect of salinity on the zooplankton community in the Pearl River Estuary. *Journal of Ocean University of China*, 19(6), 1389-1398. <https://doi.org/10.1007/s11802-020-4449-6>

Topics of Limnological Research in Mexico, coordinated by Alfredo Pérez Morales, was published in Dirección General de Publicaciones of the Universidad de Colima, avenida Universidad 333, Colima, Colima, México, www.ucol.mx. The edition was completed in December 2025. The Arial family was used for typesetting. Non Periodical Editorial Program: Eréndira Cortés Ventura. Cover Design: Adriana Minerva Vázquez Chávez. English Proofreader: Yul Ceballos. Interior Design: José Luis Ramírez Moreno and Leticia Bermúdez Aceves.



This book takes a significant step in showcasing the relevance of limnology to our survival. Freshwater habitats, though they cover less than 1 % of the Earth's surface, are home to a substantial portion of the world's biodiversity—at least 10 % of all known species. Freshwater habitats and the biodiversity they support are under threat. Moreover, our survival depends on access to high-quality freshwater. This book not only highlights the beauty of limnology and the scientific methods used to study it, but it also draws attention to the major causes of biodiversity loss in freshwater ecosystems. It shows all readers what it means to deal with inland waters as a scientist interested in understanding ecosystems and protecting them.

ISBN: 978-968-9733-13-3

9 789689 733133



UNIVERSIDAD DE COLIMA